

AD-A077 227 SRI INTERNATIONAL MENLO PARK CA F/G 7/5
MECHANISMS OF MULTIPHOTON DISSOCIATION OF MOLECULAR IONS.(U)
NOV 79 M J COGGIOLA , J R PETERSON N00014-76-C-1035
NL

UNCLASSIFIED

| OF |
AD
A077227



END
DATE
FILMED
12-79
DDC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

**READ INSTRUCTIONS
BEFORE COMPLETING FORM**

| REPORT DOCUMENTATION PAGE | | | READ INSTRUCTIONS BEFORE COMPLETING FORM | |
|---|-----------------------|--|--|--|
| 1. REPORT NUMBER | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER | | |
| 4. TITLE (and subtitle) Mechanisms of Multiphoton Dissociation of Molecular Ions. | | 5. TYPE OF REPORT & PERIOD COVERED Summary Questionnaire November 1978-November 1979 | | |
| 7. AUTHOR(s) Michael J. Coggiola and James R. Peterson | | 8. PERFORMING ORG. REPORT NUMBER | | |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS SRI International 333 Ravenswood Avenue Menlo Park, CA 94025 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12 4 | | |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research 800 North Quincy Street Arlington, VA 22217 | | 12. REPORT DATE NOVEMBER 1979 | | |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) LFVEI | | 13. NUMBER OF PAGES 30 | | |
| 16. DISTRIBUTION STATEMENT (of this Report) | | 15. SECURITY CLASS. (of this report) Unclassified | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | 18a. DECLASSIFICATION/DOWNGRADING SCHEDULE R D C REF ID: A62461 E NOV 23 1979 | | |
| 18. SUPPLEMENTARY NOTES | | | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multiphoton dissociation, polyatomic ions, infrared absorption, dissociation dynamics. <i>This report discusses</i> | | | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The dissociation of molecular ions resulting from infrared photon absorption by highly vibrationally excited molecular ions will be studied. Photofragments will be identified and their dissociative lifetime determined where possible. Measurements will also be made of the fragment translational energy distribution result to date on ions such as CF_3I^+ , CF_3Br^+ , and C_3F_6^+ show that a strong correlation exists between this single photon dissociation process and the final step of multiphoton dissociation. Preliminary analysis indicates that the dissociation dynamics follow a simple statistical mechanism. | | | | |

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

10. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Multiphoton dissociation, polyatomic ions, infrared absorption, dissociation dynamics.

This report discusses

20. **ABSTRACT** (Continue on reverse side if necessary and identify by block number)
 The dissociation of molecular ions resulting from infrared photon absorption by highly vibrationally excited molecular ions will be studied. Photofragments will be identified and their dissociative lifetime determined where possible. Measurements will also be made of the fragment translational energy distribution results to date on ions such as CF_3I^+ , CF_3Br^+ , and C_3F_6^+ show that a strong correlation exists between this single photon dissociation process and the final step of multiphoton dissociation. Preliminary analysis indicates that the dissociation dynamics follow a simple statistical mechanism.

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE

~~UNCLASSIFIED~~

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

November 1979

SUMMARY QUESTIONNAIRE

ONR Contract N00014-76-C-1035

Mechanisms of Multiphoton Dissociation of Molecular Ions

1. Contract Description

The dissociation of molecular ions resulting from infrared photon absorption will be studied. Photofragments will be identified and their dissociative lifetime determined where possible. Measurements will also be made of the fragment translational energy distributions.

2. Scientific Problem

In contrast to the well-characterized mechanisms of infrared multiphoton absorption and dissociation in neutral molecules, the MPD of molecular ions has only recently been demonstrated and its mechanism and dynamics remain largely unknown. The present study will provide valuable information on fundamental questions of the absorption spectra of molecular ions, the dissociative lifetime of highly excited ions, the dynamical details of the energy redistribution within such ions and the degree to which MPD in ions is a statistical thermodynamic process.

In general, this work will probe, in detail, the nature and properties of highly excited molecular ions, and their absorption of infrared radiation. This will result in a clearer characterization of the quasi-continuum of molecules and of the final step in MPD.

3. Scientific and Technical Approach

Highly vibrationally excited positive ions are formed in an electron impact source and focused into a highly collimated beam. This beam is merged over a distance of ~ 35 cm with the output from a line tunable CW CO₂ laser. Ion fragments resulting from the single photon photodissociation of ions whose initial internal energy was within $\hbar\nu$ of the dissociation limit are identified. High resolution electrostatic energy analysis of the fragments yields the center of mass kinetic energy distribution. Dissociative lifetimes are determined by measuring the photofragment signal over two different path lengths. Photodissociation cross sections are measured as a function of CO₂ laser wavelength to obtain information on the vibrational frequencies of the ions.

79 11 16 020

4. Progress

The infrared single photodissociation cross section was measured for a wide variety of molecular ions, such as CF_3I^+ , CF_3Br^+ , $\text{C}_2\text{F}_5\text{I}^+$, C_3F_6^+ and C_6F_6^+ . Many of the ions studied showed very strong resonant absorption behavior which has been identified with the ν_1 symmetric C-F stretch mode of the ion. The dissociation pathway has been confirmed in each case to follow the least endothermic channel, as found in neutral MPD. Dissociative lifetimes were also measured for several ions, and found to correlate qualitatively with molecular complexity. The absorption and subsequent dissociation process observed here is interpreted corresponding directly with the final step in MPD of ions, and hence provides new information on the details of that phenomenon. An estimation of the photodissociation cross section of CF_3I^+ indicates that approximately 0.4% of the ions in the beam have sufficient internal excitation to participate in the observed process. The measured lifetimes and kinetic energy distributions of the fragments correlate with a simple statistical picture of the redistribution of internal energy within an ion.

5. Publications

A paper entitled, "Infrared Photodissociation of Molecular Ions", by M. J. Coggiola, P. C. Cosby and J. R. Peterson has been submitted for publication in the Journal of Chemical Physics.

6. Extenuating Circumstances

None

7. Funds

We anticipate having no funds remaining at the end of the current contract period.

8. Graduate Students and Postdoctoral Personnel

No graduate students or postdoctoral personnel participated in this work.

9. Graduate Student Degrees

None

10. Other Government Support

See accompanying Table.

| | | | |
|------------------------|--|--------------------|---------------|
| Accession for GRAPH | NTIS TAB | DDC Unannounced | Justification |
| By Dist | Distribution Availability or Availability Special | | |
| R | | | |

CURRENT AND PROJECTED RESEARCH SUPPORT

| SRI Project No. | Title | Sponsor | Expiration | Annual Budget | Percentage of Professional Time per Project | | | | | |
|-----------------|--|---------|------------|---------------|---|----------|--------|----------|----------|--------|
| | | | | | Peterson | Coggioia | Hodges | Peterson | Coggioia | Hodges |
| 7155 | Collision Processes Involving Metastable Atoms and Molecules | NSF | 5/30/80 | \$78,000 | | | | 5% | 25% | 50% |
| - | Electron Capture and Loss Cross Sections and Angular Scattering of D ⁺ , O ⁻ in Cs | DOE | 11/1/80 | \$89,000 | | | | 24% | 25% | 32% |
| 6772 | Properties of Excited States of Molecular Ions | ARO | 9/11/80 | \$50,000 | | | | Peterson | Coggioia | Hodges |
| 7433/2 | Ion Photofragment Spectroscopy | AFOSR | 5/30/80 | \$60,000 | | | | 6% | 34% | 10% |
| This Proposal | | | | \$55,000 | | | | 20% | 8% | 12% |